


## Alvetex® Scaffold 384-well Plate: Quick Start Protocol

### Description

Reinnervate's Alvetex Scaffold 384-well plate is comprised of a black plate, clear plastic base, with Alvetex Scaffold at the bottom of each well. The Alvetex Scaffold has been heat welded to the base of the wells in a process which does not alter its physical structure. Cells growing in 3D are exposed to culture medium from above only and predominantly reside in the top portion of scaffold. As with 2D culture, the use of coating reagents to enhance cell growth can be considered (e.g. ECM proteins, poly-d-lysine).

Reinnervate's Alvetex Scaffold 384-well plate technology is reproducible and compatible with a variety of in vitro cell viability assays and offers the advantage of assay multiplexing, which may not be possible when using alternative cancer cell biology models, such as soft agar clonogenic assays.



**FIGURE 1:** Alvetex Scaffold in 384-well plate format.

### Notes before starting

- Reinnervate's Alvetex Scaffold 384-well plate is supplied gamma irradiated and sterile until opened.
- Prior to use, the sample wells must be rehydrated by the addition of 70 % ethanol (~75  $\mu$ l).
- The ethanol solution must then be removed by aspirating, and samples wells washed at least once with PBS (~75-100  $\mu$ L) and then once with the appropriate culture media (~75-100  $\mu$ l), or alternatively 2 x media washes (~75-100  $\mu$ l).

**Note:** The PBS and media washes must be carried out immediately after aspiration of the ethanol solution to avoid the scaffold drying out.

- The 384-well plate can now be used immediately for seeding of cells or placed in a humidified incubator, with wells containing culture media, until the cells are ready for seeding.

## Alvetex® Scaffold 384-well Plate: Quick Start Protocol

### Seeding of cells on Alvetex Scaffold 384-well plate

For the best results it is recommended to optimise cell seeding density in the first instance. We have previously demonstrated that between 1,000 and 10,000 cells can be seeded per well in the final culture volume (e.g. 75-100 µl per well). This will depend on the nature of the cell type and its appropriate growth conditions. Typically seeding cells in the range of 3,000 to 5,000 cells per well is a good starting point. Note that a 1-hour incubation at room temperature prior to culture at 37 °C may improve the uniformity of cell coverage on each scaffold. Media changes can be performed every 1-2 days but this is dependent on cell type and growth characteristics. Media changes should ideally be performed every  $\geq 2$  days to reduce any temperature variability that may be caused by repeated removal of the plate from the incubator.

**NB. Temperature variations, edge effects and stacking:** As with conventional 2D culture in the 384 well format, plates either cultured in or removed from an incubator at 37°C in tall stacks, will take longer to equilibrate to the surrounding temperature than plates arranged separately. In addition, it is well known that inefficient temperature equilibration can result in temperature gradient effects between wells in the centre and wells at the edges of a plate. Plate stacking may exacerbate such cross-plate temperature gradients as well as causing differential gradient patterns between plates within the same stack. This may affect assay performance and cell migration towards the edge of the wells when seeding. This is a generic feature that occurs in all standard micro-titre plates and is not due to the presence of Alvetex Scaffold.

Given that pipetting errors may also occur when manually removing the product from a 384-well plate it is recommended that the 384-well plate is used on an automated liquid handling platform due to small volumes dispensed.


### Example optimisation of cell seeding for Alvetex Scaffold 384-well plate format

- Use standard 2D detachment methods and count cells using Trypan Blue and haemocytometer.
- PC-3 cells were seeded at the following cell density for evaluation: 1,000, 2,000, 5,000 and 10,000 cells per well in a final volume of 75 µL. Cells were grown for periods of 1, 4 and 7 days and their growth and viability assessed by standard MTS cell viability assay.
- 10 µL of MTS reagent was added directly to each sample well containing 75 µl culture media.
- 384 well plates were incubated for one hour at 37 °C / 5 % CO<sub>2</sub> for 60 minutes in the dark, and then 60 µL of product was removed and the absorbance measured at 490 nm in a standard 384-well plate.
- After 24 hours of culture, a linear MTS assay response with increasing cell number was observed (Figure 1).
- The viability curve shows similar growth patterns for 1,000, 2,000 and 5,000 cells seeded per well. In this example, a cell seeding density of 10,000 cells per well, resulted in slower growth compared to the lower cell seeding numbers which may indicate that that the optimal numbers of cells per well was exceeded (Figure 2).


From these observations, the optimal growth conditions for PC-3 cells on the 384-well plate is recommended to be between 1,000 to 5,000 cells per well for up to four days 3D culture.

# Alvetex® Scaffold 384-well Plate: Quick Start Protocol

Example data derived from Alvetex® Scaffold 384-well plates



**FIGURE 2:** Cell number standard curve for the PC-3 cell line using the MTS cell viability assay. PC-3 cells were seeded at a range of densities onto Alvetex Scaffold 384-well plate format and cultured for 24 hours prior to analysis: linear increase in absorbance with increasing cell seeding density was observed for up to 10,000 cells per well. Data from multiple sample replicates are shown (n = 12, mean ± SD).



**FIGURE 3:** Cell number standard curves for the PC-3 cells using the MTS cell viability assay. PC-3 cells were seeded at a range of densities onto Alvetex Scaffold 384-well plate format and cultured for up to 7 days hours prior to analysis. Data represent mean ± SD, n=12.

For research use only. Not for use in diagnostic procedures.



## Alvetex® Scaffold 384-well Plate: Quick Start Protocol

### Technical Support

To obtain the latest technical support information, please visit [www.reinnervate.com](http://www.reinnervate.com). You can have 24-hour access to the following information:

- Access to telephone and fax information for order support
  - Search through the Reinnervate protocol video gallery
  - Search product spec sheets, protocols, applications notes and product support docs
  - Obtain information on customer training and workshops
  - Submit a question directly to Technical Support at [tech.support@reinnervate.com](mailto:tech.support@reinnervate.com)
- 

### Ordering Information

For the latest products and services available through Reinnervate, please visit [www.reinnervate.com](http://www.reinnervate.com).

#### Reinnervate

NETPark Incubator, Thomas Wright Way,  
Sedgefield, Co. Durham, TS21 3FD  
UNITED KINGDOM

T: +44 (0) 1740 625 266

F: +44 (0) 1740 625 251

E: [info@reinnervate.com](mailto:info@reinnervate.com)

---

### Trademarks

Unless otherwise noted, all trademarks are the property of Reinnervate Ltd.  
©2015 Reinnervate Ltd. All rights reserved.